




Max. Marks: 300

Date: 21.10.2022

JB 2 MR BATCH
PHYSICS : PART TEST SET-B
Topic: FLT

- Two vectors \vec{a} and \vec{b} inclined at an angle θ have a resultant \vec{c} which makes an angle β with \vec{a} . If the directions of \vec{a} and \vec{b} are interchanged, then the resultant will have the same
 - magnitude
 - direction
 - magnitude as well as direction
 - neither magnitude nor direction
- The sum of two forces acting at a point is 16 N. If the resultant force is 8 N and its direction is perpendicular to minimum force then the forces are
 - 6 N and 10 N
 - 8 N and 8 N
 - 4 N and 12 N
 - 2 N and 14 N
- Rain is falling with speed $12\sqrt{2}$ m/s at an angle of 45° with vertical line. A man in a glider going at a speed of u at angle of 37° with respect to ground. Find the speed of glider so that rain appears to him falling vertically. Consider motion of glider and rain drops in same vertical plane.




- 15 m/s
- 30 m/s
- 10 m/s
- 25 m/s


Space for Rough Work




4. A body of mass 1 kg lies on smooth inclined plane. The block of mass m is given force $F = 10$ N horizontally as shown. The magnitude of net normal reaction on the block is:



- (a) $10\sqrt{2}$ N (b) $\frac{10}{\sqrt{2}}$ N (c) 10 N (d) None of these
5. A body of mass 2.0 kg is placed on a smooth horizontal surface. Two forces $F_1 = 20$ N and $F_2 = N$ are acting on the body in directions making angles of 30° and 60° to the surface. The reaction of the surface on the body will be



- (a) 20 N (b) 25 N (c) 5 N (d) zero
6. M is a fixed wedge. Masses m_1 and m_2 are connected by a light string. The wedge is smooth and the pulley is smooth and fixed. $m_1 = 10$ kg and $m_2 = 7.5$ kg. When m_2 is just released, the distance it will travel in 2 seconds is




- (a) 2.8 m (b) 7.5 m (c) 4.0 m (d) 6.0 m


Space for Rough Work




7. With what acceleration should the box of figure be moving up so that the block of mass M exerts a force $7Mg/4$ on the floor of the box?



- (a) $g/4$ (b) $g/2$ (c) $3g/4$ (d) $4g$
8. In the given diagram, with what force must the man pull the rope to hold the plank in position? Mass of the man is 80 kg. Neglect the weights of plank, rope and pulley. Take $g = 10 \text{ ms}^{-2}$.



- (a) 200 N (b) 300 N (c) 600 N (d) 150 N
9. Three blocks 1, 2 and 3 are arranged as shown in the figure. The velocities of the blocks v_1 , v_2 and v_3 are shown in the figure. What is the relationship between v_1 , v_2 and v_3 ?



- (a) $2v_1 + v_2 = v_3$ (b) $v_1 + v_2 = v_3$ (c) $v_1 + 2v_2 = v_3$ (d) None of these


Space for Rough Work



10. A block of mass 2 kg rests on a rough inclined plane making an angle of 30° with the horizontal. The coefficient of static friction between the block and the plane is 0.7. The frictional force on the block is

(a) 10 N (b) $7\sqrt{3}$ N (c) $10 \times \sqrt{3}$ N (d) 7 N

11. In the arrangement shown in the figure [$\sin 37^\circ = 3/5$]



(a) direction of force of friction is up the plane (b) the magnitude of force of friction is zero
(c) the tension in the string is 20 N (d) magnitude of force of friction is 56 N


12. The work done by external agent in stretching a spring of force constant $k = 100$ N/cm from deformation $x_1 = 10$ to deformation $x_2 = 20$ cm.

(a) -150 J (b) 50 J (c) 150 J (d) None of these

13. An elastic spring of unstretched length L and force constant K is stretched by amount x . It is further stretched by another length y . The work done in the second stretching is

(a) $\frac{1}{2} Ky^2$ (b) $\frac{1}{2} K(x^2 + y^2)$ (c) $\frac{1}{2} K(x + y)^2$ (d) $\frac{1}{2} Ky(2x + y)$

14. A toy car of mass 5 kg moves up a ramp under the influence of force F plotted against displacement x . The maximum height attained is given by




(a) 20 m (b) 15 m (c) 11 m (d) 5 m


Space for Rough Work



15. A slab S of mass m is released from a height h_0 from the top of a spring of force constant k . The maximum compression x of the spring is given by the equation



- (a) $mgh_0 = \frac{1}{2} kx^2$ (b) $mg(h_0 - x) = \frac{1}{2} kx^2$
 (c) $mgh_0 = \frac{1}{2} k(h_0 + x)^2$ (d) $mg(h_0 + x) = \frac{1}{2} kx^2$
16. A particle travels 10 m in first 5 sec and 10 m in next 3 sec. Assuming constant acceleration what is the distance travelled in next 2 sec
- (a) 8.3 m (b) 9.3 m (c) 10.3 m (d) None of above
17. A car moving at 160 km/h when passes the mark-A, driver applies brake and reduces its speed uniformly to 40 km/h at mark-C. The marks are spaced at equal distances along the road as shown below.
 At which part of the track the car has instantaneous speed of 100 km/h? Neglect the size of the car.




- (a) At mark-B (b) Between mark-A and mark-B
 (c) between mark-B and mark-C (d) insufficient information to decide
18. A particle is moving along a straight line with constant acceleration. At the end of tenth second its velocity becomes 20 m/s and in tenth second it travels a distance of 10 m. Then the acceleration of the particle will be
- (a) 10 m/s^2 (b) 20 m/s^2 (c) $\frac{1}{5} \text{ m/s}^2$ (d) 3.8 m/s^2


Space for Rough Work



19. Two cars are moving in the same direction with the same speed 30 km/hr. They are separated by a distance of 5 km, the speed of a car moving in the opposite direction if it meets these two cars at an interval of 4 minutes, will be
 (a) 40 km/hr (b) 45 km/hr (c) 30 km/hr (d) 15 km/hr
20. An elevator, in which a man is standing, is moving upward with a constant acceleration of 2 m/s^2 . At some instant when speed of elevator is 10 m/s, the man drops a coin from a height of 1.5 m. Find the time taken by the coin to reach the floor.
 (a) $\frac{1}{\sqrt{3}}$ sec (b) $\frac{1}{2}$ sec (c) $\frac{1}{\sqrt{2}}$ sec (d) 1 sec
21. The velocity-time graph of a body moving in a straight line is shown in the figure. The displacement and distance travelled by the body in 6 sec are respectively



- (a) 8 m, 16 m (b) 16 m, 8 m (c) 16 m, 16 m (d) 8 m, 8 m
22. The path of a projectile in the absence of air drag is shown in the figure by dotted line. If the air resistance is not ignored then which one of the path shown in the figure is appropriate for the projectile




- (a) B (b) A (c) D (d) C

Space for Rough Work



23. A stone is projected from the ground with velocity 50 m/s at an angle of 30° . It crosses a wall after 3 sec. How far beyond the wall the stone will strike the ground ($g = 10 \text{ m/sec}^2$)?
- (a) 90.2 m (b) 89.6 m (c) 86.6 m (d) 70.2 m
24. A particle starts moving with velocity 10 m/s in a straight line under an acceleration varying linearly with time. Its velocity time graph is as shown in figure. Its velocity is maximum at $t = 3$ sec. The time (in sec) when the particle stops is ($\tan 37^\circ = 3/4$)



- (a) 6 (b) 8 (c) 7 (d) 5
25. When two vectors of magnitudes P and Q are inclined at an angle θ the magnitude of their resultant $2P$. When the inclination is changed to $180 - \theta$ the magnitude of the resultant is halved. Find the ratio of P to Q.
- (a) $\sqrt{2} : \sqrt{3}$ (b) $1 : \sqrt{3}$ (c) $1 : \sqrt{2}$ (d) $\sqrt{3} : \sqrt{2}$

Space for Rough Work

Date: 21.10.2022

JB 2 MR BATCH
CHEMISTRY : PART TEST SET-B
Topic: FLT

26. Mendeleev predicted the physical and chemical properties of the elements. Eka – boron, Eka – silicon and Eka – aluminium. The elements were discovered respectively as
(a) Ge, Sc, Ga (b) Sc, Ge, Ga (c) Al, Ga, Ge (d) Ge, Al, Ga
27. Lothar Meyer obtained the curve for the known elements by plotting their atomic volumes against
(a) atomic numbers (b) atomic masses (c) densities (d) ionization energies
28. The correct order of the size of C, N, P and S is
(a) $N < C < P < S$ (b) $C < N < P < S$ (c) $C < N < S < P$ (d) $N < C < S < P$
29. (A), (B) and (C) are elements in the third short period. Oxide of (A) is ionic, that of (B) is amphoteric and that of (C) is a giant molecule. Then (A), (B) and (C) have atomic number in the order
(a) $(C) < (B) < (A)$ (b) $(A) < (B) < (C)$ (c) $(A) < (C) < (B)$ (d) $(B) < (A) < (C)$
30. How many Cs atoms can be converted to Cs^+ ions by 1 joule energy if IE_1 for Cs is 376 KJ mol^{-1}
(a) 1.6×10^{18} (b) 1.6×10^{10} (c) 5.8×10^{14} (d) 5.8×10^{25}
31. Equivalent weight of H_2O_2 is
(a) 17 (b) 34 (c) 68 (d) 18
32. 2 g of a metal oxide on strong heating gave 112 ml of O_2 gas at STP. Equivalent of the metal is
(a) 56 (b) 92 (c) 112 (d) 50
33. One mole of chlorine combines with certain weight of metal giving 111 g of its chloride. The same amount of metal can displace 2gm hydrogen from an acid. The equivalent weight of metal is....
(a) 40 (b) 20 (c) 80 (d) 10
34. For which of the following compound equivalent weight is equal to molecular weight
(a) H_2SO_4 (b) H_3PO_2 (c) H_3PO_4 (d) H_3PO_3

Space for Rough Work



35. The atomic number of element Unq is :
 (a) 102 (b) 103 (c) 104 (d) 105
36. The temperature at which RMS velocity of SO_2 molecules is half that of He molecules at 300 K
 (a) 150 (b) 600 K (c) 900 K (d) 1200 K
37. Which is correct with respect to 'p' orbitals
 (a) They are spherical (b) They have a strong directional character
 (c) They are five fold degenerate (d) They have no directional character
38. The maximum number of electrons in an atom with $l = 2$ and $n = 3$ is
 (a) 2 (b) 6 (c) 12 (d) 10
39. Containers A and B have the same gas. Pressure, volume and temperature of A are all twice those of B, then the ratio of number of molecules of A to B is
 (a) 1 : 2 (b) 4 : 1 (c) 1 : 4 (d) 2 : 1
40. The following electronic transition corresponds to the shortest wave length ($n = \text{no. of orbit}$)
 (a) $n_5 \rightarrow n_1$ (b) $n_5 \rightarrow n_3$ (c) $n_5 \rightarrow n_2$ (d) $n_5 \rightarrow n_4$
41. What is the wave number of yellow radiation having wavelength 5800 \AA (in cm^{-1})
 (a) 5.17×10^{14} (b) 1.7×10^4 (c) 4×10^{14} (d) 3.4×10^{14}
42. Photoelectron emission is observed from a surface for frequencies ν_1 and ν_2 and of the incident radiation ($\nu_1 > \nu_2$). If the maximum kinetic energies of the photoelectrons in the two cases are in the ratio 1 : K then the threshold frequency ν_0 is given by
 (a) $\frac{\nu_2 - \nu_1}{K - 1}$ (b) $\frac{k\nu_1 - \nu_2}{K - 1}$ (c) $\frac{k\nu_2 - \nu_1}{K - 1}$ (d) $\frac{\nu_2 - \nu_1}{K}$
43. The maximum number of electrons that can be accommodated in a 3d subshell is
 (a) 2 (b) 10 (c) 6 (d) 14

Space for Rough Work



44. Which of the following statements concerning sunlight is false?
- (a) It is a form of energy (b) It cannot be deflected by a magnet
(c) It consists of photons of same energy (d) It is a part of electromagnetic spectrum
45. Number of waves produced by an electron in one complete revolution in n^{th} orbit is:
- (a) n (b) n^2 (c) $(n + 1)$ (d) $(2n + 1)$
46. One would expect proton to have very large
- (a) Ionization potential (b) Radius
(c) Charge (d) Hydration energy
47. Atoms have a mass of the order
- (a) 10^{-26} kg (b) 10^{-15} kg (c) 10^{-26} g (d) 10^{-15} g
48. The electronic velocity in fourth Bohr orbit of hydrogen is V . The velocity of electron in the first Bohr orbit would be :
- (a) $4V$ (b) $16V$ (c) $V/4$ (d) $V/16$
49. At STP 5.6 L of a gas weigh 60 g. The vapour density of gas is:
- (a) 60 (b) 120 (c) 30 (d) 240
50. 9 g of trivalent metal combines with 8 g of oxygen. The atomic mass of the metal is
- (a) 9 (b) 18 (c) 27 (d) 36

Space for Rough Work

JB 2 MR BATCH
MATHEMATICS : PART TEST SET-B
Topic: FLT

51. A question paper is divided into two parts A and B and each part contains 5 questions. The number of ways in which a candidate can answer 6 questions selecting at least two questions from each part is
(a) 200 (b) 100 (c) 80 (d) None of these
52. The value of $e^{(\log_{18} \cot 1^\circ + \log_{10} \cot 2^\circ + \dots + \log_{10} \cot 89^\circ)}$ is
(a) 0 (b) $\frac{1}{e}$ (c) 1 (d) e
53. $\sin^2 5^\circ + \sin^2 10^\circ + \sin^2 15^\circ + \dots + \sin^2 85^\circ + \sin^2 90^\circ =$
(a) $9\frac{1}{2}$ (b) 9 (c) 8 (d) 7
54. If $\log_{\cos x} \sin x = 0.5$, when $0 < x < 90^\circ$, then the value of $\cos x =$
(a) $\frac{\sqrt{5} + 1}{4}$ (b) $\frac{\sqrt{5} - 1}{4}$ (c) $\frac{\sqrt{5} - 1}{2}$ (d) $\frac{\sqrt{5} + 1}{2}$
55. If $\frac{\cos^4 \alpha}{\cos^2 \beta} + \frac{\sin^4 \alpha}{\sin^2 \beta} = 1$ then the value of $\frac{\cos^4 \beta}{\cos^2 \alpha} + \frac{\sin^4 \beta}{\sin^2 \alpha}$ is equal to
(a) 0 (b) 1 (c) 2 (d) 4
56. The total number of selections of at most n things from $(2n + 1)$ different things is 63. Then the value of n is
(a) 4 (b) 3 (c) 2 (d) None of these
57. A total numbers of words which can be formed out of the letters a, b, c, d, e, f taken 3 together such that each word contains at least one vowel is
(a) 48 (b) 96 (c) 72 (d) None of these

Space for Rough Work



58. A father with 8 children takes them 3 at a time to zoological gardens, as often as he can without taking the same 3 children together more than once. The number of times each child will go to the garden is
 (a) 112 (b) 21 (c) 56 (d) None of these
59. A dictionary is printed consisting of 7 lettered words only than can be made with a letter of the word CRICKET. If the words are printed at the alphabetical order, as in an ordinary dictionary, then the number of word before the word CRICKET is
 (a) 481 (b) 530 (c) 531 (d) 480
60. If a, b, c are three natural numbers in A.P. and $a + b + c = 21$, then the possible number of values of the ordered triplet (a, b, c) is
 (a) 13 (b) 15 (c) 14 (d) None of these
61. The product of the roots of the equation $x^2 - 4mx + 3e^{2\log m} - 4 = 0$ is 8, then its roots will be real when m equals
 (a) $\pm\sqrt{2}$ (b) ± 2 (c) $\sqrt{2}$ (d) 1
62. The number of solution of the equation $\sin(a^x) + \cos(a^x) = a^x + a^{-x}$ is $a > 0$
 (a) 3 (b) 1 (c) 2 (d) 0
63. If α, β are roots of the equation $ax^2 + 3x + 2 = 0$ ($a < 0$) then $\frac{\alpha^2}{\beta} + \frac{\beta^2}{\alpha}$ is greater than
 (a) 2 (b) 1 (c) 0 (d) None of these
64. The product of real roots of the equation $|2x + 3|^2 - 3|2x + 3| + 2 = 0$ is
 (a) 2 (b) $\frac{5}{4}$ (c) $\frac{5}{2}$ (d) 5

Space for Rough Work



65. Two students while solving a quadratic equation in x with leading coefficient '1' one copied the constant term incorrectly and got the roots 3 and 2. The other copied the constant term correctly -6 . The correct roots are
 (a) $-1, 6$ (b) $-6, -1$ (c) $-2, 3$ (d) $-3, 2$
66. If α and β are the roots of equation $8x^2 - 3x + 27 = 0$ then $\left(\frac{\alpha^2}{\beta}\right)^{1/3} + \left(\frac{\beta^2}{\alpha}\right)^{1/3}$ is equal to
 (a) $\frac{1}{4}$ (b) $\frac{1}{3}$ (c) $\frac{7}{2}$ (d) $\frac{1}{16}$
67. If α and β are the roots of $x^2 - 2x + 4 = 0$ then $\alpha^5 + \beta^5$ is equal to
 (a) 16 (b) 32 (c) 64 (d) None of these
68. In a geometric progression consisting of positive terms each term equals the sum of the next two term. Then the common ratio of this progression equals
 (a) $\frac{1}{2}(\sqrt{5} - 1)$ (b) $\frac{1}{2}\sqrt{5}$ (c) $\sqrt{5}$ (d) $\frac{1}{\sqrt{2}}(1 + \sqrt{5})$
69. $1^3 - 2^3 + 3^3 - 4^3 + \dots + 9^3 =$
 (a) -475 (b) 425 (c) 475 (d) -425
70. If $\log_0(3^{1+x} + 2)$, $\log_3(4 + 3^x - 1)$ are in A.P., then x equals
 (a) $1 - \log_3 4$ (b) $\log_4 3$ (c) $\log_3 4$ (d) $1 - \log_4 3$
71. The sum of the series $\frac{1}{2} + \frac{3}{4} + \frac{7}{8} + \frac{15}{16} + \dots$ upto n terms is
 (a) $n - 1 + \frac{1}{2^n}$ (b) $n + \frac{1}{2^n}$ (c) $2n + \frac{1}{2^n}$ (d) $n + 1 + \frac{1}{2^n}$
72. If $x^2 + 9y^2 + 25z^2 = xyz\left(\frac{15}{x} + \frac{5}{y} + \frac{3}{z}\right)$, then x, y and z are in
 (a) A.P. (b) G.P. (c) A.G.P. (d) H.P.

Space for Rough Work



BJNP

Learning with the Speed of Mumbai and the Tradition of Kota



73. $2^{1/4}, 4^{1/8}, 8^{1/16} \dots$ up to ∞ is equal to

- (a) 1 (b) 2 (c) $\frac{3}{2}$ (d) None of these

74. The sum of n terms of the series

$$\frac{1}{1 + \sqrt{3}} + \frac{1}{\sqrt{3} + \sqrt{5}} + \frac{1}{\sqrt{5} + \sqrt{7}} + \dots \text{ is}$$

- (a) $\sqrt{2n+1}$ (b) $\frac{1}{2}\sqrt{2n+1}$ (c) $\sqrt{2n+1} - 1$ (d) $\frac{1}{2}(\sqrt{2n+1} - 1)$

75. The value of ${}^{40}C_{31} + \sum_{j=0}^{10} {}^{40+j}C_{10+j}$ is equal to

- (a) $2 \cdot {}^{45}C_{15}$ (b) ${}^{51}C_{20}$ (c) $2 \cdot {}^{50}C_{20}$ (d) None of these

* * * * *

Space for Rough Work



Max. Marks: 300

Date: 21.10.2022

JB 2 MR BATCH
PHYSICS : PART TEST SET-B ANSWER KEY
Topic: FLT

1.	(a)	2.	(a)	3.	(a)	4.	(a)	5.	(d)
6.	(a)	7.	(c)	8.	(a)	9.	(d)	10.	(a)
11.	(a)	12.	(c)	13.	(d)	14.	(c)	15.	(d)
16.	(a)	17.	(c)	18.	(b)	19.	(b)	20.	(b)
21.	(a)	22.	(a)	23.	(c)	24.	(c)	25.	(a)

CHEMISTRY : PART TEST SET-B ANSWER KEY
Topic: FLT

26.	(b)	27.	(b)	28.	(d)	29.	(b)	30.	(a)
31.	(a)	32.	(b)	33.	(b)	34.	(b)	35.	(c)
36.	(d)	37.	(b)	38.	(d)	39.	(d)	40.	(a)
41.	(b)	42.	(b)	43.	(b)	44.	(c)	45.	(a)
46.	(d)	47.	(a)	48.	(a)	49.	(b)	50.	(c)

MATHEMATICS : PART TEST SET-B ANSWER KEY
Topic: FLT

51.	(a)	52.	(c)	53.	(a)	54.	(c)	55.	(b)
56.	(d)	57.	(b)	58.	(b)	59.	(b)	60.	(a)
61.	(b)	62.	(d)	63.	(d)	64.	(c)	65.	(a)
66.	(a)	67.	(b)	68.	(a)	69.	(b)	70.	(a)
71.	(a)	72.	(d)	73.	(b)	74.	(d)	75.	(b)